

g
14
No 31
Dec. 5 1826
S. W. 8th and Walnut

A Dissertation
on
Hydrocephalus Acutus

By Paper Read^d March 5th 1826
W. L. A.
Luc E. Nicholson

of
Virginia

Decem. 2nd 1826

[illegible]

To the Medical Faculty of the
University of Pennsylvania the
following dissertation is respectfully
submitted, for the degree of Doctor
of Medicine, by the Author: —

[Faint, illegible handwriting on the main page of the manuscript.]

[Faint handwriting on the right margin of the adjacent page, partially visible.]

Hydrocephalus Acutus. —

The diversity and contrariety of opinion entertained by medical writers on this subject together with the equivocal nature of its symptoms renders it a source of no inconsiderable embarrassment to the young and inexperienced student to write a correct dissertation on this disease. —

Hydrocephalus was divided by ancient Nosologists into, External and Internal; the former is distinguished, by an accumulation of water between the scalp and cranium the latter by a collection of water between the membranes or in the ventricles. —

Hydrocephalus Externus, is of very rare occurrence, but it does sometimes happen, and when it does, it should be regarded only

as Ror...

It is now

Water o

I have s

inangun

This

sent to c

years of

the few

by confid

stances

more suff

by suffor

by suffor

a deepie

complex

prine i

But m

dispropo

as Amaraea of the scalp says Dr. Chapman.

It is now divided by systematic writers into Acute and Chronic; The Acute species I have selected for the subject of my inaugural dissertation. —

This disease is said to be chiefly incident to children between two and six years of age, rarely occurring later than the fourteenth year; that it is not exclusively confined to children, we have numerous instances on record to prove. Adults being more subject to its attacks than is generally supposed. Those adults who most frequently suffer from this disease, are those of a dropical diathesis; pale and sallow complexion; and much affected with pains in the head. —

But when we reflect on the great disproportion between the head and other

part
great
and
is in
fierce
and
exposed
are
saying
frequency
and
brother
female
to
mean
to
uniform
Hgd

parts of the body in young subjects; the great vascularity, and laxity of the brain producing an undue determination of blood to that viscus: by which the ~~disposition~~ to effusion is increased, together with the many and diversified accidents to which the violent amusements of children constantly expose them, (which act as remote causes) we cannot hesitate for a moment in saying that children should be the most frequent sufferers by this dreadful malady; and those of a scrupulous and Ricketty Mother most subject to its attacks, and females more so than males according to Morgagni. But from subsequent researches on this disease, we are disposed to think this opinion vague and unfounded. —

Hydrocephalus Acutus generally

various
 forms of pe
 reason m
 and heet,
 but tongue
 and extrac
 ing of the
 great thin
 have been
 written in
 14
 My love
 desired as a
 the decision
 symptomatic
 found ble
 of them alto
 20
 You photo
 remembrance
 him in

4
commences with all the ordinary symptoms of fever. As languor, loss of appetite, nausea with occasional vomiting, increased heat, hurried respiration, flushed face, foul tongue, pain in the head back and extremities, accelerated pulse, throbbing of the temporal arteries, attended with great thirst, exacerbations always taking place towards the evening, and becoming milder in the morning. —

Hydrocephalus acutus may be considered as exhibiting three stages, though the division is very arbitrary, for the symptoms of the different stages will be found blended together or one or more of them altogether wanting. —

Symptoms of the first stage; usually commences with languor, drowsiness, pain in the head and back of the

with his
face his
mouth
at the time
had his
mouth
was in
pursued
pursued
much inter
becoming
the more,
my very re
the same in
which was
than my
two days.
of others
which was

neck, hot dry skin, foul tongue, flushed
 face hurried respiration, quick and
 sometimes irregular pulse, dilatation
 or contraction of one or both pupils, a
 hard dry or phlegmy cough, obstinate
 constipation of the bowels, though the
 reverse may occur, when the stools are
 procured by medicine, they are of a
 greenish appearance, sleep is very
 much interrupted, frequently starting and
 screaming as if much alarmed, picking
 the nose grinding the teeth, and becom-
 ing very restless, with exacerbations in
 the evening and the symptoms becoming
 milder in the morning -

These symptoms continuing for a
 few days only; when they are succeeded
 by others denoting the second stage
 which are characterized by the patient

the least
 part of
 day of the
 hour on
 the same
 to the ex-
 hibition
 those, for
 the longer
 by the
 from the
 consistent
 of a day
 appear to
 with the
 the whole
 be assured
 results are

8
increasing less sensible to the pain in
the head and other parts of the system,
pneumaturae slow and irregular
ity of the pulse, the pupils more dilated
than in the first stage, strabismus,
the vomiting ceases except when raised
to the erect posture, respiration becomes
laborous, the skin hot and dry, great
thirst, frequent sighing and groaning,
the tongue very white and ~~moist~~ dry,
the bowels remain very much con-
fined, the faces vary both in colour and
consistence, they are most commonly
of a clay colour, in small quantities and
appear to have an oily substance mixed
with them, The patient at this time
takes whatever is offered him either
food or medicine: and swallows it with great
avidity, and he appears to be quite inco-

[illegible]

breast and stomach. We are now pre-
sented with the symptoms of the third
stage, which are still more violent and
certainly indicate a fatal termination.

There are widely dilated pupils, the
eyelids half closed, with the axis of one or
both turned in toward the nose, the eye is
sometimes suffused with blood, the pulse
returns again to the febrile state of exite-
ment, becoming so weak and quick that it is
impossible to count it, the flushing of the
face is more frequent and succeeded by a
deadly paleness, rolling the head from side to
side on the pillow, throwing the hands about
involuntarily, the respiration very laborious
and performed very slowly, difficult deglutition
the features are shrunk, involuntary discha-
rge of the feces and urine, the urine is in
large quantity perfectly clear and limpid, cold

Hammer

seems

The

justified

consequence

many of

When

very rare

partial

substitute

a small

the room

learn

it would

stretch

find a

because

be very

the desire

slaming sweat, discharges upon now imperious
nerves, succeeded by convulsions and death.

The emaciation (as observed by a distin-
guished writer) is singularly rapid what is
commonly the effect of an atrophy of
many months taking place in a few days -

When the progress of the disease has been
very rapid, it is not uncommon for the
violent symptoms to subside, inducing an
expectation of a speedy recovery, but there is
a most threatening and fatal calm, showing
the result of exhaustion. - The vessels of the
brain being previously in a high degree
of excitement are in this way relaxed; and the
blood unobstructed, for a short time, the effused
fluid acting as a reexciting cause and the
disease returns with redoubled violence, -

We may consider the case ^{and by} then as a
most desperate or nearly so. -

...
fatal, &
on the
not to m
that the
of them
the case
and to
guaranty
But I
very few
Hydroc
the d
it seems
rays are
night and
may be
In 1800
more or less

Effusion is regarded by some writers as certainly fatal, as they deny the existence of abscesses in the brain. But the phenomena of growth not to mention other facts, sufficiently prove, that there must exist in every part of the system: for an abscess is as necessary in the composition of a living body as a blood vessel each being indispensable to the execution of its primary functions.

But it must be acknowledged that they act very feebly and incompletely in the Hydrocephalic affections. —

The duration of this disease is very uncertain it sometimes proves fatal in three or four days and at others it is protracted to seven or eight weeks but the average term of its duration may be stated at twenty one days. —

In examinations, the ventricles are found more or less distended, with fluid, varying in

guarantee

is to be

that a

list of

Richard

in the

other

to the

in the

is a

in the

a note

the reg

very

appear

on the

The

in the

state of

quantity, from one to six or eight ounces, it is sometimes as thin as water, and at others thick andropy. The veins are engorged with blood of a dark gummy colour, adhesions and thickening of the membranes of the brain tumours of different sizes, situated in either the substance of the brain or attached to the membranes, &c substances of the brain pink and white the abdominal viscera also exhibit marks of disease as inflammation of the Liver, Stomach, spleen or intestines. - It has often happened that the symptoms of Hydrocephalus have been very strongly marked, when no marked appearance could be discovered in the brain on dissection.

The remote or predisposing causes of this disease appear to consist in a certain state of inactivity and fullness of constitution

herd
author
of the
supper
perpet
a slave
prova
which
in order
but now
her power
supper
expans
enrich
How can
maternal
H
placed
heart

hereditary predisposition and the scrofulous
system. The exciting cause is the irritation
of the stomach and bowels, as produced in various
suppression, of lunar spots or discharges from
scrofulous ulcers about the neck and head; but
in those upon the head, tumours with. The
consequence, and all former diseases of the skin
that last, have a great tendency to terminate
in inflammation or effusion in the brain
and more particularly the meninges. Last but not
least, enough chronic catarrhs, toothache
suppression of the nasal hemorrhages, stoma-
chic affections, and diarrhoea. Seceding of certain
eruptions from the surface, and in fact all
those causes which tend to produce inflam-
mation in general. —

The proximate cause of this disease is
placed by Dr. Chapman, than which it
would be vain to look for higher authority.

man

Unrepe-

the effec

ed propo

cause of

action of

that are a

a disorde

of its de

for 1722

comatous

slight on

p. 100. 101.

2^d Nov

only as an

along with

It appears

first day

proceed

is an increased and altered action in the
 vessels of the brain, the disease being mostly
 the effect. Furthermore, says the latter, such
 is the effect. Though I place the proximate
 cause of Hydropcephalus ventris in a morbid
 action of the brain, I am not the less persuaded
 that in a majority of cases it commences in
 a diseased state of the stomach or some
 of its dependencies. —

Dr Beddoe believes it to belong to inflammation. —
 Dr Withering observes that
 slight inflammation and congestion are the
 precursors to the aqueous accumulation.

Dr Rush thinks that it should be regarded
 only as an effect of a primary inflammation
 a congestion of blood in the brain.

It appears says he that the disease in its
 first stage is the effect of causes, which
 produce a less degree of that inflammation

which
stage is
constituted
the form
flammar
Athens
to have
of the ab
cause of
the abse
secondary
go th
of every
a state of
retires
hard than
which sta
ability.

For

which constitutes Phrenitis and its second stage is a less degree of that effusion which constitutes serous apoplexy in adults.

The former partakes of the chronic inflammation of Dr Cullen, and the Arthritic inflammation of Dr Brown.

Dr Darwin supposes inactivity and torpor of the absorbent vessels of the brain to be the cause of this disease, though the torpor of the absorbent vessels may often exist as a secondary effect.

Dr Whist observes that the immediate cause of every kind of dropsy is the same; such a state of the parts as makes the exhalant arteries throw out a greater quantity of fluid than the absorbents can take up which state he considers as consisting in solubility.

From the authorities as quoted above

Small
and the
and the
lately
same
power
the year
in the
passed
author
compar
The
Lectures
lyphon
n. 11 m
Notes of
in path
young
to the

14
It will appear that it arises from congestion
and liquid inflammation of the brain
and sometimes from a general debility and
laxity of that organ. The former opinion
can be inclined to think the most plausible
from the appearances on dissection and
the general mode of treatment pursued
in this disease. That it does sometimes
proceed from the latter we have sufficient
authority to prove, but those cases are
comparatively few.

The diseases which Hydrocephalus
acute most resembles, are some of the
typhoid states of fever, apoplexy, &c. &c.
It may be distinguished from the typhoid
states of fever, by frequent remissions, by
idiopathic fever not being common in
young subjects while there are some subjects
to the attacks of Hydrocephalus then results.

It differs
with the
and we
attacks
we can
between
where
is trans-
greater
by page
of the
Pro-
stage of
the sec-
in the
therefor-
deserve
definite
in part

It differs from apoplexy, by being attended with fever, more common to children and not as sudden and violent in its attacks. There are no symptoms by which we can correctly ascertain the difference between this disease, and worms, except where worms are present, the abdomen is tumid and tense. We may arrive at a greater degree of certainty in the diagnosis by paying strict attention to the history of the case from its commencement.

Reveries, may take place from the first stage of this disease; very rarely from the second, and never from the third. In the progress of this disease when delirium stupor, unduly dilated pupils, strabismus, deafness, laborious respiration, difficult deglutition loss of sight, weak quick and irregular pulse, the patient cannot be raised

to the
order
the
and
perfect
sions
fallen
The
search
all of
in
marginal
turn of
water
the
the
when
gradual
various

to the erect posture without its producing violent pain in the head, and sickness at the stomach, involuntary discharges of feces and urine the urine in large quantity perfectly clear and limpid, attended by convulsions the disease will certainly terminate fatally in a very short time.

The disease ^{is} most frequently seated in the Arachnoid Membrane of the brain, though all of the membranes exhibit marks of disease or affection. It sometimes depends upon congestion in the blood vessels, or inflammation of the substance of the brain itself.

Water may continue in the ventricles of the brain many months without producing the symptoms of Hydrocephalus Acutus, when the accumulation has been very gradual. and on other times all the pathognomonic symptoms of Hydrocephalus Acutus

are part
of disease
with
to the
maternal
taken to
to soldiers
the under
had, 1800
copious
the
from
tempora
The gen
determin
the viol
thulep
passion
will be of

17

are present without discovering any marks
of disease in the bowels, or accumulation of
water within the ventricles.

In the treatment of the first or inflam-
matory stage of this disease the first indi-
cation to be attended to is to prevent oppression
to sustain vascular action and diminish
the undue determination of blood to the
head, which is best effected by bloodletting,
copious purging, occasional vomiting,
blisters &c. to derive the most benefit
from bloodletting we should open the
temporal artery or jugular vein.

The quantity to be drawn can only be
determined by the age of the patient and
the violence of the symptoms present.
Unless we bleed intelligently, a perceptible im-
provement be made on the circulation it
will be of but little benefit, if any at all.

the
sitting
handy
D
from
proceed
renew
treated
that
C
give an
found
seam
ed at
vacu
have
situated
Ant
pregn

(If the indispensable necessity, of blood
letting (say a distinguished writer) seem
hardly, express myself too strongly.

Dr. Charcote was in the habit of bleeding
from the jugular vein till fainting was
produced, the patient being placed in the
recumbent posture, of similar cases
treated in this way Dr. Charcote states
that ~~vital~~ recovery.

Immediately after bleeding we should
give an active purgative of calomel
combined with coloc or rhubarb, gamboge
scammony & aloes. which should be repeat-
ed at short intervals, until copious
evacuations are produced. After the bowels
have been thoroughly emptied of their
vitiated contents, we should resort to the
antimonial preparations in small and
frequently repeated doses so as to keep

a com
aptitude
the he
and pr
be kept
clear
lots of
on the
on him
water
conserv
often as
directed
been p
Local
the in, b
unbided
the pass
Albiste

a constant nausea at the stomach to ~~expel~~ the determination of blood to the head to relax the capillaries of the surface and promote diaphoresis. The bowels must be kept in a laxative ^{state} by the use of Rhubarb alone or the mineral salts in small doses.

Cold applications to the head are very useful, in the early stages of this disease, cloths wet in vinegar and water, ether and water, or ice under slown, and applied to the head is the most convenient mode of using it. Changing them as often as they become warm, a stream of water directed immediately on the head has often been productive of the happiest effects.

Local bleeding is likewise indispensable when the inflammatory symptoms have not entirely subsided, cups are preferable to leeches from the promptness of their action.

Blisters should not be neglected at this stage.

of the
List of the
hope of
my dear
the creature
can with
sufficient
at 17. 10. 11
in the
the exam
proposed
against 10
to effect
to. India
from
stage na
have labo
and so
into 10

of the complaint, they should discontinue the use of the most sweets and saccharine and a discharge kept up from them for several days. Glysters may perhaps, *The Ceratium cantharidis*, or the *Ceratium sabine* may be used for this purpose either of which is capable of producing a sufficient degree of excitement. In Menstrual excitement has likewise been recommended in the dropping of the uterus. Caustic applied to the cervix is said by some writers to be preferable to blisters, and it is a more powerful stimulant and more permanent in its effects it may perfectly be more useful. The use of seton is preferred for that purpose. When the symptoms of the second stage have commenced indicating effusion to have taken place bleeding will be expected and some times prove vigorous. The head should be bled and a blister be applied externally

untill
 and not
 they
 the law
 will str
 give ear
 me long
 do I an
 ought to
 negotiat
 advance
 your leg
 and with
 to Amer
 the Mex
 and larg
 can man
 than so

over its surface; which should remain on
until the supplicative process has commenced
and it will be of but little benefit.

Purgatives should likewise be used to rid
the bowels of their acid contents; calomel
with its ordinary adjuncts answer this purpose
best. Emetics combined with purgatives
are very valuable in this stage of the disorder.

Dr. Cassmichael Smith recommends very
highly the squill combined with calomel,
scitatis is sometimes used but ~~with~~ little
advantage, the neutral salts scammony,
gamboge, aloes and rhubarb, are sometimes
used with good effects, but they are all inferior
to mercury. After effusion has taken place
the medicine should be given in as large
and frequently repeated doses as the patient
can bear; it should be applied also by inunction
so as to excite salivation as speedily as

proposed
the riding
took off
which c
and the
It be
the pica
then
a soap
bless, a
sinks,
into our
remedies
of health
Be
that after
is lost his

possible. Dr. Lobson and Percival says that the relief was decided as soon as the Mercury took effect. We believe it to be the only remedy which can cure the disease after effusion and this very often fails.

Dr. Cheyne tells us he succeeded in curing the disease with the James' powder.

When the patient is convalescent he should be supported with, barrow root, sago, tapioca, ellies, light broths &c mild and stimulating drinks, wine with proper tonics and more note exercise in the open air. These are the remedies most conducive to the restoration of health. —

But in conclusion it must be admitted that after effusion has taken place there is but little to be expected from our remedies.

